



8/3/2021

Lilly and Incyte's baricitinib reduced deaths among patients with COVID-19 receiving invasive mechanical ventilation

- New data from Phase 3 COV-BARRIER sub-study indicates one death prevented for every six baricitinib-treated patients on mechanical ventilation or ECMO compared to placebo

- Data showed 46% risk reduction in mortality by Day 28 and 44% risk reduction in mortality by Day 60

INDIANAPOLIS, Aug. 3, 2021 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) and Incyte (NASDAQ:INCY) announced today results from an additional cohort of 101 adult patients from the COV-BARRIER trial. In this sub-study, patients with COVID-19 on mechanical ventilation or extracorporeal membrane oxygenation (ECMO) who received baricitinib plus standard of care were 46 percent less likely to die by Day 28 compared to patients who received placebo plus standard of care (nominal p-value=0.0296; hazard ratio [HR] [95% CI] = 0.54 [0.31, 0.96]; analysis not adjusted for multiplicity). The cumulative proportion of patients who died by Day 28 was 39.2 percent (n/N: 20/51) in the baricitinib arm versus 58 percent in the placebo arm (n/N: 29/50). Similar mortality benefit was observed by Day 60 (HR [96% CI] = 0.56 [0.33, 0.97]) with a cumulative proportion of death of 45.1 percent (n/N: 23/51) for baricitinib compared to 62.0 percent for placebo (n/N: 31/50). These findings are consistent with the reduction in mortality observed in the overall COV-BARRIER patient population.

"As additional data from COV-BARRIER become available, it is increasingly evident that treatment with baricitinib may help prevent death in some of the most critically ill COVID-19 patients and that baricitinib represents an important treatment option for this vulnerable group of patients in this constantly evolving pandemic," said E. Wesley Ely, M.D., M.P.H., professor of medicine and co-director of the Critical Illness, Brain Dysfunction, and Survivorship (CIBS) Center at Vanderbilt University Medical Center and co-principal investigator of COV-BARRIER.

By Day 28, the frequency of adverse events, serious adverse events and serious infections were similar in the baricitinib group (88%, 50% and 44%, respectively) compared to placebo (95.9%, 71.4% and 53.1%, respectively). Venous thromboembolic events were reported in 6 percent of patients treated with baricitinib and 6.1 percent of patients treated with placebo. No new safety signals were identified.

"In the interest of public health and safety, it remains a priority to provide healthcare professionals with as much



information as possible about treatment options that may help improve outcomes for patients with severe disease," said Ilya Yuffa, senior vice president and president of Lilly Bio-Medicines. "These new data add to the growing body of evidence demonstrating the important role baricitinib has and may continue to play for certain hospitalized patients with COVID-19."

Lilly intends to publish detailed results from this additional sub-study in a peer-reviewed journal and present the findings at a medical meeting in the coming months. These new data from the COV-BARRIER sub-study will also be shared with regulatory authorities in the U.S., European Union and other geographies.

On July 28, 2021, the U.S. Food and Drug Administration (FDA) **broadened** the Emergency Use Authorization (EUA) for baricitinib to allow for treatment with or without remdesivir. The EUA provides for the use of baricitinib for treatment of COVID-19 in hospitalized adults and pediatric patients two years of age or older requiring supplemental oxygen, non-invasive or invasive mechanical ventilation or ECMO. For more information about the authorized use of baricitinib in COVID-19 and mandatory requirements of the EUA, please see the **FDA Letter of Authorization, Fact Sheet for Healthcare Providers** and Fact Sheet for Patients, Parents and Caregivers (**English**) (**Spanish**).

Baricitinib is an oral JAK inhibitor discovered by Incyte and licensed to Lilly.

Authorized Use Under the EUA and Important Safety Information for baricitinib (in the United States) for COVID-19

Baricitinib is authorized for use under an Emergency Use Authorization (EUA) for treatment of COVID-19 in hospitalized adults and pediatric patients 2 years of age or older requiring supplemental oxygen, non-invasive or invasive mechanical ventilation, or extracorporeal membrane oxygenation (ECMO).

Baricitinib has not been approved for the treatment of COVID-19, but has been authorized for emergency use by the FDA. Baricitinib is authorized under an EUA only for the duration of the declaration that circumstances exist justifying the authorization of the EUA of baricitinib under Section 564(b)(1) of the Act, 21 U.S.C. § 360bbb-3(b)(1), unless the declaration is terminated or authorization revoked sooner.

For more information about the authorized use of baricitinib in COVID-19 and mandatory requirements of the EUA, please see the **FDA Letter of Authorization, Fact Sheet for Healthcare Providers** and Fact Sheet for Patients, Parents and Caregivers (**English**) (**Spanish**).

Important Safety Information about baricitinib for COVID-19

The following provides essential safety information on the unapproved use of baricitinib under the Emergency Use

Authorization.

Warnings

Serious Infections: There is limited information regarding use of baricitinib in patients with COVID-19 and concomitant active serious infections.

Serious infections have occurred in patients receiving baricitinib. Avoid the use of baricitinib with known active tuberculosis. Consider if the potential benefits outweigh the potential risks of baricitinib treatment in patients with active serious infections other than COVID-19 or chronic/recurrent infections.

Thrombosis: In hospitalized patients with COVID-19, prophylaxis for venous thromboembolism is recommended unless contraindicated. If clinical features of deep vein thrombosis or pulmonary embolism occur, patients should be evaluated promptly and treated appropriately.

Abnormal Laboratory Values: There is limited information regarding use of baricitinib in patients with COVID-19 and any of the following clinical findings: absolute neutrophil count (ANC) <1000 cells/mm³, absolute lymphocyte count (ALC) <200 cells/mm³, and hemoglobin <8 g/dL.

Evaluate estimated glomerular filtration rate (eGFR), liver enzymes, and complete blood count at baseline and thereafter according to local patient management practice. Monitor closely when treating patients with abnormal baseline and post-baseline laboratory values. Follow dose adjustments as recommended in the Fact Sheet for Healthcare Providers for patients with abnormal renal, hematological and hepatic laboratory values. Manage patients according to routine clinical guidelines.

Vaccinations: Avoid use of live vaccines with baricitinib.

Hypersensitivity: If a serious hypersensitivity occurs, discontinue baricitinib while evaluating the potential causes of the reaction.

See **Warnings and Precautions** in the FDA-approved full **Prescribing Information** and **Medication Guide** for additional information on risks associated with longer-term treatment with baricitinib.

Serious Side Effects

Serious venous thrombosis, including pulmonary embolism, and serious infections have been observed in COVID-19 patients treated with baricitinib and are known adverse drug reactions of baricitinib.

Adverse Reactions

In the COVID-19 clinical trials, adverse drug reactions in the safety population occurring in $\geq 1\%$ of patients treated with baricitinib were alanine aminotransferase (ALT) ≥ 3 x upper limit of normal (ULN) (18.0%), aspartate aminotransferase (AST) ≥ 3 x ULN (11.5%), thrombocytosis $>600,000$ cells/mm³ (8.2%), creatine phosphokinase (CPK) >5 x ULN (3.7%), neutropenia <1000 cells/mm³ (2.2%), deep vein thrombosis (1.5%), pulmonary embolism (1.4%), and urinary tract infection (1.3%).

Use in Specific Populations

Pregnancy: Baricitinib should be used during pregnancy only if the potential benefit justifies the potential risk for the mother and the fetus.

Renal Impairment: There are limited data for baricitinib in patients with severe renal impairment. Baricitinib is not recommended for patients who are on dialysis, have end-stage renal disease, or have acute kidney injury.

Hepatic Impairment: Baricitinib has not been studied in patients with severe hepatic impairment. Baricitinib should only be used in patients with severe hepatic impairment if the potential benefit outweighs the potential risk.

Please see **Fact Sheet for Healthcare Providers** and **Fact Sheet for Patients, Parents and Caregivers (English)** or **Fact Sheet for Patients, Parents and Caregivers (Spanish)** .

BC HCP EUA ISI 28JUL2021

About OLUMIANT® (baricitinib)

OLUMIANT, a once-daily, oral JAK inhibitor was discovered by Incyte and licensed to Lilly. It is approved in the U.S. and more than 75 countries as a treatment for adults with moderate to severe rheumatoid arthritis. It is also approved for the treatment of certain hospitalized patients with COVID-19 in Japan. The U.S. FDA-approved labeling for OLUMIANT includes a Boxed Warning for Serious Infections, Malignancy, and Thrombosis. See the full Prescribing Information [here](#).

In December 2009, Lilly and Incyte announced an exclusive worldwide license and collaboration agreement for the development and commercialization of baricitinib and certain follow-on compounds for patients with inflammatory and autoimmune diseases.

About COV-BARRIER Sub-Study

COV-BARRIER sub-study was a randomized, double-blind, placebo-controlled trial, initiated in December 2020, to assess the efficacy and safety of baricitinib versus placebo when added to standard of care which included corticosteroids (86% of the patients) in patients hospitalized with COVID-19 requiring invasive mechanical ventilation or ECMO at baseline. A total of 101 patients were randomized to baricitinib or placebo with 51 receiving baricitinib and 50 receiving placebo. The pre-specified endpoints include 28-days mortality (and 60-day) and the number of ventilator-free days. All analyses are exploratory and no multiplicity adjustment was applied.

COV-BARRIER was a global, randomized, double-blind, placebo-controlled study of hospitalized patients comparing baricitinib 4 mg once daily plus standard of care versus placebo plus standard of care. Patients could remain on background standard of care, as defined per local guidelines, including antimalarials, antivirals, corticosteroids, and/or azithromycin. The most frequently used therapies were corticosteroids (79% of patients, mostly dexamethasone) and remdesivir (19% of patients). While the composite primary endpoint of COV-BARRIER, which was defined as a difference in the estimated proportion of participants progressing to non-invasive ventilation including high flow oxygen or invasive mechanical ventilation (including ECMO) or death by Day 28, did not meet statistical significance, baricitinib-treated patients (27.8%) were less likely than those receiving standard of care (30.5%) to progress to ventilation or death (odds ratio [OR]: 0.85; 95% CI: 0.67, 1.08; p=0.180). A pre-specified key secondary endpoint showed baricitinib, in addition to standard of care, meaningfully reduced the risk of death by 39 percent by Day 28 when compared to standard of care alone (n/N: 62/764 [8.1%] baricitinib, 101/761 [13.3%] placebo; [estimated difference in Day 28 probability of mortality = -4.9% (95% CI: -8.0%, -1.9%); hazard ratio [HR] = 0.56 (95% CI: 0.41, 0.77)]. No new safety signals were identified. The study findings from COV-BARRIER have been submitted to a peer-reviewed journal for future print publication.

About Lilly's COVID-19 Efforts

Lilly is bringing the full force of its scientific and medical expertise to attack the coronavirus pandemic around the world. Existing Lilly medicines are being studied to understand their potential in treating complications of COVID-19, and the company is collaborating with partner companies to discover and develop novel antibody treatments for COVID-19. Click [here](#) for resources related to Lilly's COVID-19 efforts.

About Eli Lilly and Company

Lilly is a global health care leader that unites caring with discovery to create medicines that make life better for people around the world. We were founded more than a century ago by a man committed to creating high-quality medicines that meet real needs, and today we remain true to that mission in all our work. Across the globe, Lilly employees work to discover and bring life-changing medicines to those who need them, improve the

understanding and management of disease, and give back to communities through philanthropy and volunteerism. To learn more about Lilly, please visit us at [lilly.com](https://www.lilly.com) and [lilly.com/newsroom](https://www.lilly.com/newsroom). P-LLY

About Incyte

Incyte is a Wilmington, Delaware-based, global biopharmaceutical company focused on finding solutions for serious unmet medical needs through the discovery, development and commercialization of proprietary therapeutics. For additional information on Incyte, please visit [Incyte.com](https://www.incyte.com) and follow [@Incyte](https://twitter.com/Incyte).

Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about OLUMIANT (baricitinib) as a potential treatment for patients with COVID-19 and reflects Lilly's and Incyte's current beliefs and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of research, development and commercialization. Among other things, there can be no guarantee that planned or ongoing studies will be completed as planned, that future study results will be consistent with the results to date, that OLUMIANT will receive additional regulatory approvals or authorizations or be commercially successful, that OLUMIANT will be safe and effective as a treatment for COVID-19, or that we can provide an adequate supply of OLUMIANT in all circumstances. For further discussion of these and other risks and uncertainties, see Lilly's and Incyte's most recent respective Form 10-K, Form 10-Q, and Form 8-K filings with the United States Securities and Exchange Commission. Except as required by law, Lilly and Incyte undertake no duty to update forward-looking statements to reflect events after the date of this release.

Refer to: Rachel Sorvig; sorvig_rachel@lilly.com; +1-317-607-7507 (Lilly media)
Kevin Hern; hern_kevin_r@lilly.com; +1-317-277-1838 (Lilly investors)
Catalina Loveman; cloveman@incyte.com; +1-302-498-6171 (Incyte media)
Christine Chiou; cchiou@incyte.com; +1-302-274-4773 (Incyte investors)

View original content to download multimedia: <https://www.prnewswire.com/news-releases/lilly-and-incytes-baricitinib-reduced-deaths-among-patients-with-covid-19-receiving-invasive-mechanical-ventilation-301346651.html>

SOURCE Eli Lilly and Company

